

HESSIAN EQUATIONS WITH ELEMENTARY SYMMETRIC FUNCTIONS

HONGJIE DONG

ABSTRACT. We consider the Dirichlet problem for two types of degenerate elliptic Hessian equations. New results about solvability of the equations in the $C^{1,1}$ space are provided.

1. INTRODUCTION

This article is closely related to [11], [7] and [15]. A second-order partial differential equation is called *Hessian* equation if it is of the form

$$F(u_{xx}) = f,$$

where (u_{xx}) is the Hessian matrix of u and $F(w)$ only depends on the eigenvalues of the symmetric matrix w .

Here we are concerned with the Dirichlet problem for two types of degenerate Hessian equations:

$$P_m(u_{xx}) = \sum_{k=0}^{m-1} (l_k^+)^{m-k}(x) P_k(u_{xx}), \quad (1.1)$$

$$P_m(u_{xx}) = g^{m-1}, \quad (1.2)$$

where $P_k(u_{xx})$ is the k th elementary symmetric polynomial of eigenvalues of the matrix u_{xx} , and l_k, g are in $C^{1,1}$. The second equation is called m -Hessian equation, which becomes the *Monge-Ampère* equation when $m = d$. Equation (1.2) is non-degenerate if $g > 0$ and is degenerate if g can vanish at some points. Many authors have studied the Hessian equations and especially the Monge-Ampère equation. The solvability of *non-degenerate* equations is proved by establishing the $C^{2,\alpha}$ estimate of the solutions and using the method of continuity (see [1], [13] and [14]). For non-degenerate equations, we also refer the

Date: February 2, 2008.

1991 *Mathematics Subject Classification.* 35J60.

Key words and phrases. Degenerate Hessian equations; Dirichlet problem; Weak solutions.

reader to the works [10], [1], [3] and [6]. It is well-known that there exists a unique admissible weak solution to *degenerate* problem (see [22]). To show the existence and uniqueness of solutions in $C^{1,1}$, it suffices to obtain a priori estimate of the second order derivatives of smooth solutions to the approximating non-degenerate equations (cf. Lemma 4.6).

For the degenerate case, a global upper bound for the second-order derivatives of admissible solutions to the Dirichlet problem of Hessian equations, and more general *Bellman* equations, was established by N.V. Krylov in a series of papers [15]-[18]. In [15] a few concrete equations similar to (1.1) and (1.2) like

$$P_m(u_{xx}) = \sum_{k=0}^{m-1} (l_k^+)^{m-k+1}(x) P_k(u_{xx}),$$

$$P_m(u_{xx}) = (l_k^+)^{m-k}(x) P_k(u_{xx}), \quad k < m,$$

and, in particular, $P_m(u_{xx}) = (g_+)^m$ as $k = 0$, are treated as the applications of the general theory.

Later, for the case of the Monge-Ampère equation

$$P_d(u_{xx}) = \det(u_{xx}) = g^{d-1},$$

the solvability in the $C^{1,1}$ space was proved in [7] by P. Guan, N.S. Trudinger and X-J. Wang with a different approach. The power $d-1$ of g was also shown to be optimal by an example in [23]. A modification of such example shows that $m-1$ is the lowest possible power of g when there exist second derivatives estimates for solutions of the m -Hessian equations (see also recent [11], where the authors did a very good survey of the literature on the Hessian equations). As pointed out in [7], the techniques there rest on the fact that the solution is convex in case $m = d$, which in general does not always hold true for m -Hessian equations. With g^{m-1} on the right-hand side, the solvability of the general degenerate m -Hessian equations in the $C^{1,1}$ space is still unknown.

The purpose of this paper is to prove the solvability of the Dirichlet problems of type (1.1) with $C^{3,1}$ boundary data, and also the Dirichlet problems for the degenerate m -Hessian equations (1.2) with *homogeneous* boundary data. Our results improve the corresponding results in [15]. Quite a few arguments in the paper are based on or follow the results in [15] and [17]. The technique we use is to reduce the Hessian equations to the elliptic Bellman's equations and then apply the general theorems on the Bellman equations, which were introduced in [15].

Owing to an observation that a certain function is *quasiconvex* (Theorem 3.4) we are able to apply this technique to show the solvability of equation (1.2).

The article is organized as follows. Our two main theorems (2.5 and 2.6) are given in the following section. Theorem 2.5 is proved in Section 3. We prove some preliminary results and give the estimates of u, u_x in Section 4 and 5. In Section 6, we use the maximum principle in a sub-domain and reduce the estimation of second order derivatives to the estimation of their values on the boundary. After that, the boundary second derivatives are estimated in Section 7 in a standard way combining with a Hopf type lemma (Lemma 5.3), and this completes the proof of Theorem 2.6.

To conclude the introduction, we explain some notation used in what follows: $\mathbb{R}^d$ is a d -dimensional Euclidean space with a fixed orthonormal basis. A typical point in $\mathbb{R}^d$ is denoted by $x = (x^1, x^2, \dots, x^d)$. As usual the summation convention over repeated indices is enforced. For any $l = (l^1, l^2, \dots, l^d) \in \mathbb{R}^d$ and any differentiable function u on $\mathbb{R}^d$, we denote $D_l u = u_{x^i} l^i$ and $D_l^2 u = u_{x^i x^j} l^i l^j$, etc.

Let $d \geq 2$, m be positive integers, $2 \leq m \leq d$. We denote by $\mathbb{S}^d$ the set of all symmetric $d \times d$ matrices, $\mathbb{R}_+$ ($\mathbb{R}_+^0$) the set of all nonnegative (strictly positive) real numbers and $\mathcal{M}_+$ ($\mathcal{M}_+^0$) the set of all nonnegative (strictly positive) symmetric $d \times d$ matrices.

Various constants are denoted by N and δ in general and the expression $N = N(\cdot)$ means that the given constant N depends only on the contents of the parentheses.

ACKNOWLEDGEMENT

The author would like to express his sincere gratitude to his advisor, N.V. Krylov, for pointing out this problem and giving many useful suggestions and comments for improvements. The author is also grateful to the anonymous referee who gave very useful comments on the first submission of the article.

2. THE SETTING AND MAIN RESULTS

Define $P_m(\lambda) = P_{m,d}(\lambda)$ as the m th elementary symmetric polynomial of the variables $\lambda = (\lambda^1, \lambda^2, \dots, \lambda^d)$. For any symmetric $d \times d$ matrix w , define $\lambda(w)$ as a vector of eigenvalues of w with arbitrary order and define $P_m(w) = P_m(\lambda(w))$. Let $C_m = C_{m,d}$ be the open connected component of the set $\{w \in \mathbb{S}^d : P_m(w) > 0\}$ which contains the identity matrix I .

Definition 2.1. Suppose $D := \{x \in \mathbb{R}^d \mid \psi(x) > 0\}$ is a smooth bounded domain with connected boundary. We say D is $m-1$ -convex if for a large number K and any point on ∂D we have

$$P_{m-1,d-1}(\kappa^1, \dots, \kappa^{d-1}) \geq 1/K, \quad (2.1)$$

where $\kappa^1, \dots, \kappa^{d-1}$ are the principle curvatures of ∂D at this point evaluated with respect to the interior normal to ∂D .

Definition 2.2. A function $u \in C^2(D)$ is called m -admissible if the Hessian matrix (u_{xx}) is in $\bar{C}_m$ for any $x \in D$. A function $u \in C^{1,1}(D)$ is called m -admissible if the Hessian matrix of the second order Sobolev derivatives (u_{xx}) is in $\bar{C}_m$ for almost any $x \in D$.

In this article, we always assume that D is an $m-1$ -convex bounded domain of class $C^{3,1}$ with connected boundary. We are concerned with the following two Dirichlet problems for the elliptic Hessian equations,

$$P_m(u_{xx}) = \sum_{k=0}^{m-1} (l_k^+)^{m-k}(x) P_k(u_{xx}) \quad (a.e.) \text{ in } D, \quad (2.2)$$

$$u = \phi \text{ on } \partial D \quad (2.3)$$

where $l_0, \dots, l_{m-1}$ are bounded real valued functions in $\mathbb{R}^d$ and $\phi \in C^{3,1}(\mathbb{R}^d)$, and the m -Hessian equation,

$$P_m(u_{xx}) = g^{m-1} \quad \text{in } D, \quad (2.4)$$

$$u = \phi \quad \text{on } \partial D, \quad (2.5)$$

where $2 \leq m \leq d$, $g \in C^1(\bar{D})$ is nonnegative, $\phi \in C^{3,1}(\bar{D})$.

Assumption 2.3. For some sufficiently large number K and any $k = 0, 1, \dots, m-1$ the functions $l_k(x) + K|x|^2$ and $g(x) + K|x|^2$ are convex on $\bar{D}$, and

$$\text{diam}\{D\} + \|g\|_{C^1(\bar{D})} + 1/\sup_{\bar{D}} |g| + \|\phi\|_{C^{3,1}(\bar{D})} + \|\psi\|_{C^{3,1}} + \|l_k\|_{C^0} \leq K,$$

$$|\psi_x| \geq 1/K \quad \text{on } \partial D.$$

$$|\nabla g(x)|^2 \leq Kg(x) \quad \text{in } D. \quad (2.6)$$

Remark 2.4. Typically, inequality (2.6) holds if $g \in C^{1,1}(\mathbb{R}^d)$ is non-negative and $\|g\|_{C^{1,1}} \leq K/2$. We note here that (2.6) is also needed in [7] as well, for example, in the proof of Lemma 2.1 there. Because of that the result of [7] does not completely cover the result in [15] about $\det(u_{xx}) = (g_+)^d$ since $(g_+)^{d/(d-1)}$ may not satisfy (2.6).

Here come our two main results.

Theorem 2.5. *Under the above assumptions, equation (2.2)-(2.3) has a unique solution $u \in C^{1,1}(\bar{D})$ characterized by the additional property that*

$$P_m(u_{x^i x^j} + t\delta_{ij}) > 0,$$

$$P_m(u_{x^i x^j} + t\delta_{ij}) > \sum_{k=0}^{m-1} (l_k^+)^{m-k}(x) P_k(u_{x^i x^j} + t\delta_{ij}) \quad (\text{a.e.}) \text{ in } D,$$

for any $t > 0$. Moreover, if $\sum_k l_k^+ > 0$ in $\bar{D}$, then $u \in C^{2,\alpha}(\bar{D})$ for an $\alpha \in (0, 1)$.

Theorem 2.6. *Under the above assumptions, there exists a unique m -admissible solution $u \in C^{1,1}(\bar{D})$ of the Dirichlet problem (2.4)-(2.5) with homogeneous boundary condition, i.e. $\phi \equiv 0$. Moreover, the solution u satisfies (2.4) almost everywhere in D and admits an estimate*

$$\|u\|_{C^{1,1}(\bar{D})} \leq N(D, d, K). \quad (2.7)$$

Assumption 2.3 and in addition we assume more g near ∂D , that is for some large K_1 ,

3. PROOF OF THEOREM 2.5

Firstly, let's restate some results, which can be found, for instance, in [15], as the following two lemmas.

Lemma 3.1. *The set C_m defined above is an open convex cone in $\mathbb{S}^d$ with vertex at the origin containing $\mathcal{M}_+^0$, and for $k = 0, 1, \dots, m-1$ we have $C_m \subset C_k$. Furthermore, the functions $(P_k/P_{k-1})(w)$, $k = 2, 3, \dots, m$, are concave in C_m .*

Lemma 3.2. *If a function $F(w)$ is convex and homogeneous of degree $-\alpha$ in a cone $C \subset \mathbb{S}^d$, then the function $l^{\alpha+1}F(w)$ is a convex function of (w, l) in $C \times \mathbb{R}_+$.*

The following lemma can be easily proved by direct calculation of the Hessian matrix.

Lemma 3.3. *For any positive number n , the function*

$$((x + y)^n - x^n)^{-1}$$

is a convex function in the open cone $\mathbb{R}_+^0 \times \mathbb{R}_+^0$.

As a corollary of Lemma 3.2 and 3.3, the function

$$H(x, y, l) := l^{n+1}/((x + y)^n - x^n)$$

is convex in $\mathbb{R}_+^0 \times \mathbb{R}_+^0 \times \mathbb{R}_+$.

The proof of Theorem 2.5 relies on the following observation.

Theorem 3.4. *We denote*

$$G(w, l) = \sum_{k=0}^{m-1} (l_k^{m-k})(P_k/P_m)(w),$$

where $l = (l_0, l_1, \dots, l_{m-1})$. Then for any $(w, l), (\tilde{w}, \tilde{l}) \in C_m \times \mathbb{R}_+^m$, we have

$$G\left(\frac{w + \tilde{w}}{2}, \frac{l + \tilde{l}}{2}\right) \leq \max(G(w, l), G(\tilde{w}, \tilde{l})).$$

That is, $G(w, l)$ is quasiconvex in $C_m \times \mathbb{R}_+^m$.

Proof. Suppose for some $c \geq 0$, the following two inequalities hold

$$\sum_{k=0}^{m-1} l_k^{m-k} P_k(w) \leq c P_m(w), \quad \sum_{k=0}^{m-1} \tilde{l}_k^{m-k} P_k(\tilde{w}) \leq c P_m(\tilde{w}).$$

Due to the homogeneity, to prove the theorem it suffices to prove the inequality

$$\sum_{k=0}^{m-1} (l_k + \tilde{l}_k)^{m-k} P_k(w + \tilde{w}) \leq c P_m(w + \tilde{w}). \quad (3.1)$$

For $k = 0, \dots, m-1$, let $\alpha_k, \tilde{\alpha}_k$ be the nonnegative numbers such that

$$(l_k + \alpha_k)^{m-k} P_k(w) = \sum_{j=0}^k l_j^{m-j} P_j(w), \quad (3.2)$$

$$(\tilde{l}_k + \tilde{\alpha}_k)^{m-k} P_k(\tilde{w}) = \sum_{j=0}^k \tilde{l}_j^{m-j} P_j(\tilde{w}). \quad (3.3)$$

Then,

$$(l_{m-1} + \alpha_{m-1}) P_{m-1}(w) \leq c P_m(w), \quad (\tilde{l}_{m-1} + \tilde{\alpha}_{m-1}) P_{m-1}(\tilde{w}) \leq c P_m(\tilde{w}).$$

Owing to the concavity of $(P_m/P_{m-1})(w)$, we get

$$\begin{aligned} c(P_m/P_{m-1})(w + \tilde{w}) &\geq c(P_m/P_{m-1})(w) + c(P_m/P_{m-1})(\tilde{w}) \\ &\geq l_{m-1} + \alpha_{m-1} + \tilde{l}_{m-1} + \tilde{\alpha}_{m-1}, \end{aligned}$$

$$c P_m(w + \tilde{w}) \geq (l_{m-1} + \alpha_{m-1} + \tilde{l}_{m-1} + \tilde{\alpha}_{m-1}) P_{m-1}(w + \tilde{w}). \quad (3.4)$$

By using (3.2) and (3.3), for $k = 1, \dots, m-1$ we have

$$\begin{aligned} ((l_k + \alpha_k)^{m-k} - l_k^{m-k}) P_k(w) &= (l_{k-1} + \alpha_{k-1})^{m-k+1} P_{k-1}(w), \\ ((\tilde{l}_k + \tilde{\alpha}_k)^{m-k} - \tilde{l}_k^{m-k}) P_k(\tilde{w}) &= (\tilde{l}_{k-1} + \tilde{\alpha}_{k-1})^{m-k+1} P_{k-1}(\tilde{w}). \end{aligned}$$

For any $\varepsilon > 0$, the equalities above imply the following

$$\begin{aligned} ((l_k + \alpha_k + 2\varepsilon)^{m-k} - (l_k + \varepsilon)^{m-k})P_k(w) &\geq (l_{k-1} + \alpha_{k-1})^{m-k+1}P_{k-1}(w), \\ ((\tilde{l}_k + \tilde{\alpha}_k + 2\varepsilon)^{m-k} - (\tilde{l}_k + \varepsilon)^{m-k})P_k(\tilde{w}) &\geq (\tilde{l}_{k-1} + \tilde{\alpha}_{k-1})^{m-k+1}P_{k-1}(\tilde{w}). \end{aligned}$$

Again, owing to the concavity of $(P_k/P_{k-1})(w)$ in C_m and the corollary of Lemma 3.3, we obtain

$$\begin{aligned} (P_k/P_{k-1})(w + \tilde{w}) &\geq (P_k/P_{k-1})(w) + (P_k/P_{k-1})(\tilde{w}) \\ &\geq \frac{(l_{k-1} + \alpha_{k-1})^{m-k+1}}{(l_k + \alpha_k + 2\varepsilon)^{m-k} - (l_k + \varepsilon)^{m-k}} + \frac{(\tilde{l}_{k-1} + \tilde{\alpha}_{k-1})^{m-k+1}}{(\tilde{l}_k + \tilde{\alpha}_k + 2\varepsilon)^{m-k} - (\tilde{l}_k + \varepsilon)^{m-k}} \\ &\geq \frac{(l_{k-1} + \alpha_{k-1} + \tilde{l}_{k-1} + \tilde{\alpha}_{k-1})^{m-k+1}}{(l_k + \alpha_k + \tilde{l}_k + \tilde{\alpha}_k + 4\varepsilon)^{m-k} - (l_k + \tilde{l}_k + 2\varepsilon)^{m-k}}. \end{aligned}$$

As a consequence,

$$\begin{aligned} ((l_k + \alpha_k + \tilde{l}_k + \tilde{\alpha}_k + 4\varepsilon)^{m-k} - (l_k + \tilde{l}_k + 2\varepsilon)^{m-k})P_k(w + \tilde{w}) \\ \geq (l_{k-1} + \alpha_{k-1} + \tilde{l}_{k-1} + \tilde{\alpha}_{k-1})^{m-k+1}P_{k-1}(w + \tilde{w}). \end{aligned}$$

Letting $\varepsilon \downarrow 0$ and taking the limit yield

$$\begin{aligned} ((l_k + \alpha_k + \tilde{l}_k + \tilde{\alpha}_k)^{m-k} - (l_k + \tilde{l}_k)^{m-k})P_k(w + \tilde{w}) \\ \geq (l_{k-1} + \alpha_{k-1} + \tilde{l}_{k-1} + \tilde{\alpha}_{k-1})^{m-k+1}P_{k-1}(w + \tilde{w}). \end{aligned} \quad (3.5)$$

Inequality (3.1) follows if we add (3.4) and (3.5) together, and the theorem is proved. $\square$

Finally, by relying on Theorem 3.4 at one point, we can essentially reproduce Krylov's approach in [15] with very few modifications (cf. Remark 5.14, 5.16 of [15]). The idea is to reduce the equation to a Bellman's equation and apply a general existence and uniqueness result on degenerate elliptic Bellman's equations proved in [18] by a probabilistic argument, or by an analytic approach in [16] and [17]. In detail, due to Theorem 3.4 we can easily get that

$$\Theta(l) := \{w \in C_m : G(w, l) < 1\}$$

is convex in l in the sense that for any $w_i \in \Theta(l_i), i = 1, 2$ we have

$$\frac{w_1 + w_2}{2} \in \Theta\left(\frac{l_1 + l_2}{2}\right).$$

Then after one reduces the equation to a Bellman's equation, the free term is semi-concave so that the regularity theory for Bellman's equations is applicable. Theorem 2.5 improves the corresponding result in [15] also from the point of view of the following remark.

Remark 3.5. The same conclusion as in Theorem 2.5 holds true if we replace (2.2) by

$$P_m(u_{xx}) = \sum_{k=0}^{m-1} f_k(l_k^+(x)) P_k(u_{xx}) \quad (a.e.) \text{ in } D,$$

for any $s \geq 0$, where $f_k : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ are continuous functions and $f_k^{1/(m-k)}$ are convex for $k = 0, 1, \dots, m-1$.

Indeed, if we denote

$$G_1(w, l) = \sum_{k=0}^{m-1} f_k(l_k) (P_k/P_m)(w),$$

then due to Theorem 3.4 and the convexity of $f_k^{1/(m-k)}$ in $\mathbb{R}_+$, for any $(w, l), (\tilde{w}, \tilde{l}) \in C_m \times \mathbb{R}_+^m$ we have

$$\begin{aligned} & \max(G_1(w, l), G_1(\tilde{w}, \tilde{l})) \\ &= \max \left\{ \sum_k f_k(l_k) (P_k/P_m)(w), \sum_k f_k(\tilde{l}_k) (P_k/P_m)(\tilde{w}) \right\} \\ &\geq \sum_k \left[\frac{1}{2} (f_k^{1/(m-k)}(l_k) + f_k^{1/(m-k)}(\tilde{l}_k)) \right]^{m-k} (P_k/P_m) \left(\frac{w + \tilde{w}}{2} \right) \\ &\geq \sum_k f_k \left(\frac{l_k + \tilde{l}_k}{2} \right) (P_k/P_m) \left(\frac{w + \tilde{w}}{2} \right) = G_1 \left(\frac{w + \tilde{w}}{2}, \frac{l + \tilde{l}}{2} \right), \end{aligned}$$

i.e. $G_1(w, l)$ is also quasiconvex in $C_m \times \mathbb{R}_+^m$. Thus our assertion follows.

4. SOME PRELIMINARY RESULTS

In what follows, we consider the following Dirichlet problem for $2 \leq m \leq d$:

$$P_m(u_{xx}) = g^{m-1} \quad \text{in } D, \tag{4.1}$$

$$u = 0 \quad \text{on } \partial D, \tag{4.2}$$

where $g \in C^{1,1}(\bar{D})$ is nonnegative and P_m is defined in Section 2. We focus on the solution u such that $(u_{xx}) \in \bar{C}_m$ for any $x \in D$.

Lemma 4.1. *For any matrix (v_{ij}) in C_m , the d by d matrix $K(v) := (P_{m,v_{ij}}(v))$ is positive definite.*

Proof. Let η be a nonzero vector in $\mathbb{R}^d$. We have

$$\eta^T K(v) \eta = \text{Tr}(K(v) \eta \eta^T) = \frac{d}{dt} P_m(v + t \eta \eta^T) \Big|_{t=0}.$$

Since the rank of $\eta\eta^T$ is one, $P_m(v + t\eta\eta^T)$ is linear in t . Thus,

$$P_m(v + t\eta\eta^T) = P_m(v) + t\eta^T K(v)\eta.$$

Also note that $\eta\eta^T \in \bar{C}_d \subset \bar{C}_m$, for any $t \geq 0$ we have

$$v + t\eta\eta^T \in \bar{C}_m, \quad P_m(v + t\eta\eta^T) \geq 0.$$

Therefore, $\eta^T K(v)\eta \geq 0$ is nonnegative. If $\eta^T K(v)\eta = 0$, for any $t > 0$ it holds that

$$P_m(v - t\eta\eta^T) = P_m(v) > 0,$$

which implies $v - t\eta\eta^T \in C_m$ and thus $v/t - \eta\eta^T \in C_m$. Letting $t \rightarrow +\infty$ yields $-\eta\eta^T \in \bar{C}_m \in \bar{C}_1$. But this is impossible because

$$P_1(-\eta\eta^T) = \text{Tr}(-\eta\eta^T) = -|\eta|^2 < 0.$$

Hence it holds that $\eta^T K(v)\eta > 0$, and the lemma is proved. $\square$

The following corollary is an immediate consequence of Lemmas 3.1 and 4.1.

Corollary 4.2. *For any $d \times d$ matrix (v_{ij}) in C_m , all the $(d-1) \times (d-1)$ submatrices obtained by deleting the k th row and k th column of (v_{ij}) ($k = 1, \dots, d$) are in $C_{m-1, d-1}$.*

Lemma 4.3. (i) *For any matrices (v_{ij}) and (w_{ij}) in C_m , we have*

$$\text{Tr}(K(v)w) = P_{m, v_{ij}}(v)w_{ij} > 0.$$

(ii) *Moreover, for any orthogonal matrix Q , it holds that*

$$K(QvQ^T) = QK(v)Q^T. \quad (4.3)$$

Proof. The key idea of the proof of part (i) is to use the properties of hyperbolic polynomials. We treat $P_m(v)$ as a homogenous polynomial of $d(d+1)/2$ variables

$$v_{11}, v_{12}, \dots, v_{1d}, v_{22}, v_{23}, \dots, v_{2d}, v_{33}, \dots, v_{3d}, \dots, v_{dd}.$$

It's known that for any $\mu \in \mathbb{R}^d$, all roots of the polynomial $P_m(\mu + t\lambda_0)$ are real, where $\lambda_0 = (1, 1, \dots, 1)$ (cf. Corollary 6.5 of [15]). Therefore, by the very definition of P_m in $\mathbb{S}^d$, for any $v \in \mathbb{S}^d$, all roots of the polynomial $P_m(v + tI_d)$ are real. Owing to Lemma 4.16 and Theorem 6.4 of [15], for any v, w in C_m we have

$$P_{m, v_{ij}}(v)w_{ij} = \frac{d}{dt} P_m(v + tw) \Big|_{t=0} > 0.$$

This proves the first part of the lemma.

To prove the second part, first notice that P_m and C_m are both invariant under orthogonal transformations. Suppose w is a symmetric matrix. We have

$$\begin{aligned} \text{Tr}(K(QvQ^T)w) &= \frac{d}{dt} P_m(QvQ^T + tw) \Big|_{t=0} = \frac{d}{dt} P_m(v + tQ^T w Q) \Big|_{t=0} \\ &= \text{Tr}(K(v)Q^T w Q) = \text{Tr}(QK(v)Q^T w). \end{aligned}$$

Since w is an arbitrary symmetric matrix, the conclusion of (ii) follows immediately. $\square$

Remark 4.4. By using the same method as in the proof of 4.3 (i), owing to Lemma 4.16 and Theorem 6.4 of [15], one can prove that for any $(v_{ij}), (w_{ij})$ in C_m and $t_0, s_0 \geq 0$ it holds that

$$\frac{\partial^2}{\partial t \partial s} P_m(sv + tw) \Big|_{(s,t)=(s_0,t_0)} \geq 0.$$

And this implies

$$P_m(v + w) \geq P_m(v) + P_m(w). \quad (4.4)$$

Lemma 4.5. *Let $v \in \bar{C}_m$ and c be a nonnegative constant. Then $P_m(v) = c$ if and only if*

$$\inf_{w \in C_m} \{a^{ij}(w)v^{ij} - m(d - m + 1)^{-1}P_m^{1-1/m}(w)P_{m-1}^{-1}(w)c^{1/m}\} = 0, \quad (4.5)$$

where

$$a^{ij}(w) = P_{m,w^{ij}}(w) / \text{Tr}(K(w)). \quad (4.6)$$

Proof. First we suppose $P_m(v) = c$. Recall that $P_m^{1/m}(v)$ is concave in C_m (see, for instance, Theorem 6.4 of [15]). We get

$$\begin{aligned} c^{1/m} &= P_m^{1/m}(v) \\ &= \inf_{w \in C_m} \{m^{-1}P_{m,w^{ij}}(w)P_m^{1/m-1}(w)(v^{ij} - w^{ij}) + P_m^{1/m}(w)\} \\ &= \inf_{w \in C_m} \{m^{-1}P_{m,w^{ij}}(w)P_m^{1/m-1}(w)v^{ij}(x)\}. \end{aligned} \quad (4.7)$$

The last equality is because P_m is a homogeneous polynomial of degree m . In case $c > 0$, because the infimum above is attained when $\omega = v$ and also because $\text{Tr}(K(v)) > 0$, equality (4.5) follows immediately. If $c = 0$, then one has $v \in \partial C_m$. Note that

$$\text{Tr}(K(v)) = (d - m + 1)P_{m-1}(v)$$

and $v + \varepsilon I \in C_m$. For any real number $\varepsilon > 0$, we have

$$\begin{aligned} a^{ij}(v + \varepsilon I)v^{ij} &= P_{m,v^{ij}}(v + \varepsilon I)v^{ij} / \text{Tr}(K(v + \varepsilon I)) \\ &= P_{m,v^{ij}}(v + \varepsilon I)(v^{ij} + \varepsilon\delta_{ij} - \varepsilon\delta_{ij}) / \text{Tr}(K(v + \varepsilon I)) \end{aligned}$$

$$= m(d - m + 1)^{-1}(P_m/P_{m-1})(v + \varepsilon I) - \varepsilon.$$

Thanks to Lemma 4.16 (ii) of [15], we get

$$\lim_{\varepsilon \downarrow 0} a^{ij}(v + \varepsilon I)v^{ij} = 0,$$

and (4.5) follows.

On the other hand, assume that (4.5) holds true. Observe that

$$P_m(v + tI) \rightarrow +\infty \quad \text{as } t \rightarrow +\infty$$

and there exists $t_0 \leq 0$ such that $v + t_0 I \in \partial C_m$ and $P_m(v + t_0 I) = 0$. By continuity one can find a real number $t_1 \geq t_0$ such that $v + t_1 I \in \bar{C}_m$ and $P_m(v + t_1 I) = c$. Due to the first part of the proof, it holds that

$$\inf_{w \in C_m} \{ \text{Tr}(a(w)(v + t_1 I)) - m(d - m + 1)^{-1} P_m^{1-1/m}(w) P_{m-1}^{-1}(w) c^{1/m} \} = 0,$$

$$\inf_{w \in C_m} \{ \text{Tr}(a(w)v) - m(d - m + 1)^{-1} P_m^{1-1/m}(w) P_{m-1}^{-1}(w) c^{1/m} \} = -t_1.$$

Therefore, from (4.5) we obtain $t_1 = 0$, and the lemma is proved. $\square$

Because of Lemma 4.5, equation (4.1) is equivalent to

$$\inf_{w \in C_m} \{ L^w u(x) + f(w, x) \} = 0, \quad (4.8)$$

where

$$L^w u(x) = a^{ij}(w) u_{x_i x_j}(x),$$

$$f(w, x) = -m(d - m + 1)^{-1} P_m^{1-1/m}(w) P_{m-1}^{-1}(w) g^{1-1/m}(x).$$

Owing to Theorem 1.1 of [22], (4.1)-(4.2) has a unique admissible weak solution $u \in C^0(D)$ in the sense that for a sequence of m -admissible functions $u_k \in C^2(D)$ we have

$$u_k \rightarrow u \text{ in } C^0(D), \quad P_m(u_{k,xx}) \rightarrow g^{m-1} \text{ in } L^1_{\text{loc}}(D).$$

The following lemma will be proved in Section 8.

Lemma 4.6. *Under Assumption 2.3, if we can establish a priori C^2 estimate of solutions to non-degenerate problems with $C^2(\bar{D}) \in g > 0$, which does not depend on the infimum of g in D , then*

- (i) *the admissible weak solution u of (4.1)-(4.2) is in $C^{1,1}(\bar{D})$.*
- (ii) *Moreover, u satisfies (4.1) almost everywhere in D .*

Due to Lemma 4.6, from now on, we always assume that g is positive on $\bar{D}$ and belongs to $C^2(\bar{D})$. In this case, it's known that $u \in C^4(D) \cap C^2(\bar{D})$. Because the infimum in (4.8) is attained when $\omega = u_{xx}$, it is easy to see that (4.8) is equivalent to a uniformly elliptic Bellman equation

$$\inf_{w \in C_m^*} \{ L^w u(x) + f(w, x) \} = 0, \quad (4.9)$$

where

$$C_m^* = \{w \in C_m | P_m(w) \geq \inf_D g^{m-1}, w \leq N(\|u\|_{C^2(\bar{D})})I\}$$

Let ξ be a unit vector in $\mathbb{R}^d$. Denote $\lambda^1(x), \dots, \lambda^d(x)$ to be the eigenvalues of the Hessian matrix $u_{xx}(x)$. Due to Lemma 3.1, we have

$$\lambda^1 + \lambda^2 + \dots + \lambda^d = \text{Tr}(u_{xx}) = \Delta u = P_1(u_{xx}) > 0. \quad (4.10)$$

We can get more than (4.10). Define Λ_2 as the open connected component of the set $\{\lambda \in \mathbb{R}^d : P_2(\lambda) > 0\}$ which contains the vector $(1, 1, \dots, 1)$. Since $(u_{xx}) \in C_m \subset C_2$, we have $\lambda := (\lambda^1, \lambda^2, \dots, \lambda^d) \in \Lambda_2$. Obviously, $\lambda_\varepsilon := (1, \varepsilon, \varepsilon, \dots, \varepsilon)$ is also in Λ_2 for any $\varepsilon > 0$. Thus by Theorem 6.4(i) in [15] with λ_ε and P_2 in place of λ_1 and Q_m respectively, we get

$$0 < \frac{\partial}{\partial t} P_2(\lambda + t\lambda_\varepsilon) = \lambda^2 + \lambda^3 + \dots + \lambda^d + O(\varepsilon).$$

By letting $\varepsilon \downarrow 0$ and taking the limit, we get

$$\lambda^2 + \lambda^3 + \dots + \lambda^d = \text{Tr}(u_{xx}) - \lambda^1 \geq 0. \quad (4.11)$$

Of course, (4.11) remains true if we replace 1 by any $i = 2, 3, \dots, d$.

If $\max_i \lambda^i(x) < 1$, due to (4.10), we immediately get an estimate of $\lambda^i(x)$, $i = 1, 2, \dots, d$. And this yields the estimate of (u_{xx}) . Therefore, in the sequel we only consider the region

$$D' := \{x \in D | \max_i \lambda^i(x) \geq 1\}.$$

Due to (4.11), in D' we have $P_1(u_{xx}) \geq 1$.

Lemma 4.7. (i) For any $x \in D'$, we have

$$g^{m-2}(x) \leq \text{Tr}(K(u_{xx}))/ (d - m + 1). \quad (4.12)$$

(ii) For any $x \in D$, we have

$$g^{m-2}(x) \leq N g^{m-3/2}(x) \leq N \text{Tr}(K(u_{xx}))/ (d - m + 1), \quad (4.13)$$

where N depends only on K .

Proof. Note that $(P_k(w)/\binom{d}{k})$ is a log-concave function of k for $k = 0, 1, \dots, m$ (cf., for instance, Corollary 6.5 [15]). So in D' we have

$$\begin{aligned} g^{m-2}(x) &= P_m^{(m-2)/(m-1)}(u_{xx}) \leq P_m^{(m-2)/(m-1)}(u_{xx}) P_1^{1/(m-1)}(u_{xx}) \\ &\leq N P_{m-1}(u_{xx}) = N \text{Tr}(K(u_{xx}))/ (d - m + 1). \end{aligned}$$

Also for any matrix $w \in C_m$,

$$P_m^{1-1/m}(w) = P_0^{1/m}(w) P_m^{(m-1)/m}(w) \leq N P_{m-1}(w). \quad (4.14)$$

Thus in D , we have

$$\begin{aligned} g^{m-3/2}(x) &\leq Ng^{(m-1)^2/m}(x) = NP_m^{(m-1)/m}(u_{xx}(x)) \\ &\leq NP_{m-1}(u_{xx}(x)) = N\text{Tr}(K(u_{xx}))/ (d-m+1). \end{aligned}$$

The lemma is proved $\square$

Due to Assumption 2.3 and (4.14), for any $w \in C_m$ we have

$$\|f(w, \cdot)\|_{C^1(\bar{D})} \leq N(K, d). \quad (4.15)$$

Near ∂D let $\psi_2 = \text{dist}(x, \partial D)$ if $x \in \bar{D}$, $\psi_2 = -\text{dist}(x, \partial D)$ if $x \notin D$. By a standard argument in [15], we can define $\psi_1 = \psi_2 - t\psi_2^2$ near ∂D with t sufficiently large and continue ψ_1 in an appropriate manner such that

$$\{x \in \mathbb{R}^d : \psi_1(x) > 0\} = D, \quad \|\psi_1\|_{C^{3,1}(\mathbb{R}^d)} \leq N(K, d)$$

and we can find $\rho = \rho(K, d) > 0$ such that for any $x \in \bar{\Delta}_\rho$, $-\psi_{1,xx}$ is in C_m , where

$$\Delta_\rho = \{x \in D : \text{dist}(x, \partial D) < \rho\}.$$

Denote Ω to be the closure of $\{K(v)/\text{Tr}(K(v)) \mid v \in C_m\}$. Obviously, Ω is a compact set. Owing to Lemma 4.3, by a compactness argument we obtain the following corollary.

Corollary 4.8. *There exist a $\delta = \delta(K, d, \rho) > 0$, such that for any $v \in \Omega$ and $x \in \bar{\Delta}_\rho$ we have $\text{Tr}(v\psi_{1,xx}(x)) < -\delta$. Especially, for any $w \in C_m$ and $x \in \bar{\Delta}_\rho$, we have $L^w\psi_1(x) < -\delta$. Moreover, for any $x \in \partial D$, $|\psi_{1x}| = 1$.*

Proof. For any $x \in \bar{\Delta}_\rho$ we can find a number $\varepsilon > 0$ such that $-\psi_{1,xx} - \varepsilon I \in C_m$. Then we have

$$\text{Tr}(v(-\psi_{1,xx} - \varepsilon I)) \geq 0, \quad \text{Tr}(v(-\psi_{1,xx})) = \varepsilon > 0.$$

Because both Ω and $\bar{\Delta}_\rho$ are compact, there exists $\delta > 0$ such that $\text{Tr}(v\psi_{1,xx}(x)) < -\delta$. The last assertion follows immediately from the definition of ψ_1 . $\square$

Let $\psi_0 = (2R+1)^2 - |x - x_0|^2$, where R is the diameter of D and x_0 is a point in D . Since $\text{Tr}(a(w)) = 1$, it holds that $L^w\psi_0 = -2$ for any $w \in C_m$. Thus Assumption 1.2 (a), (b) in [18] are satisfied and the following lemma is proved in [18], Lemma 1.1.

Lemma 4.9. *There exist $\psi \in C^{3,1}$ and $\delta = \delta(K, d) > 0$ satisfying (i) $\{x \in \mathbb{R}^d : \psi(x) > 0\} = D$, (ii) $|\psi_x| > \delta$ on ∂D , (iii) $\|\psi\|_{C^{3,1}(\mathbb{R}^d)} \leq N(K, d)$ and (iv) for any $w \in C_m$ and $x \in D$ we have $L^w\psi(x) < -\delta$.*

5. ESTIMATES OF u AND u_x

In this section, we will give some estimates of u and u_x .

Theorem 5.1. *For any $x \in D$, we have*

$$|u(x) - \phi(x)| \leq N(K, d)\psi(x).$$

Proof. Denote $\tilde{u} = u - \phi$. Since P_m is a homogeneous polynomial of degree m , we have

$$P_{m, u_{xx}^{i_{xx}j}}(u_{xx}(x))u_{xx}^{i_{xx}j}(x) = mP_m(u_{xx}(x)) = mg^{m-1}(x),$$

$$\text{Tr}(a(u_{xx}(x))\tilde{u}_{xx}) = mg^{m-1}(x)/\text{Tr}(K(u_{xx}(x))) - \text{Tr}(a(u_{xx}(x))\phi_{xx}). \quad (5.1)$$

From the positiveness of $K(u_{xx})$, equality (5.1) can be looked at as a second order elliptic equation of $\tilde{u}$. Due to (4.13), the right hand side of (5.1) is bounded by a constant depending only on K , d and m . Lemma 4.9 implies

$$\text{Tr}(a(u_{xx}(x))\psi_{xx}(x)) < -\delta.$$

After using the comparison principle, we get what we expected. $\square$

Theorem 5.2. *We have $\sup_D |u_x| \leq N$, where N depends only on K and d .*

Proof. After differentiating (4.1) in the direction ξ , we get

$$a^{ij}(u_{xx}(x))u_{(\xi)x^i x^j}(x) = \frac{m-1}{d-m+1}g_{(\xi)}(x)g^{m-2}(x)P_{m-1}^{-1}(u_{xx}(x)). \quad (5.2)$$

Owing to (2.6) and (4.13), the absolute value of the right-hand side of (5.2) is less than

$$N|g_{(\xi)}(x)|g^{m-2}(x)P_{m-1}^{-1}(u_{xx}(x)) \leq N|g^{m-3/2}P_{m-1}^{-1}(u_{xx}(x))| \leq N.$$

Again by using Lemma 4.9 and the comparison principle, we get

$$\sup_D |u_{(\xi)}| \leq N + \sup_{\partial D} |u_{(\xi)}|.$$

Upon using Lemma 5.1, we get the estimate of the first derivative on the boundary:

$$|u_{(\xi)}(x)| \leq N(K, d) \quad \forall x \text{ on } \partial D,$$

and the lemma is proved. $\square$

We also need a lower bound for the normal first order derivative $D_n u$ on the boundary ∂D .

Lemma 5.3. *Under Assumption 2.3, for any $x \in \partial D$ we have*

$$D_n u(x) \leq -\gamma < 0,$$

where n is the inner normal of ∂D at point x and γ is a constant depending only on D , K and d .

Proof. From (4.7) with $w = I_d$ we get

$$\Delta u(x) \geq N(m, d)g^{1-1/m}(x). \quad (5.3)$$

Because g is not always equal to 0 in D and $u = 0$ on ∂D , by the strong maximum principle, we get u is strictly positive in D .

Since D is a $C^{3,1}$ domain, we can find two positive numbers ε_1 and ε_2 depending only on K satisfying: (i) for any $x \in \partial D$, there is a ball B_x of radius ε_1 which is inside D and ∂B_x and ∂D are tangent at x ; (ii) for any $x \in \partial D$, if we look at x as "north poll", then the south half ball of B_x is in $D \setminus \Delta_{\varepsilon_2}$.

Next, we claim that there exists $\gamma_0 = \gamma_0(D, d, K) > 0$ such that

$$u(x) \leq -\gamma_0 \quad \text{in } D \setminus \Delta_{\varepsilon_2}.$$

We prove this by contradiction. If this is not true, we can find two sequences $x_j \in \bar{D}$, $g_j \in C^2(\bar{D})$ such that

$$\|g_j\|_{C^1(\bar{D})} \leq K, \quad g_j(x_j) \geq 1/K, \quad (5.4)$$

and also u_j solves (4.1)-(4.2) with g_j in place of g such that

$$\sup_{D \setminus \Delta_{\varepsilon_2}} u_j \rightarrow 0, \quad \text{as } j \rightarrow +\infty. \quad (5.5)$$

Because $\bar{D}$ is compact, after passing to a subsequence, we may assume x_j converges to a point $x_0 \in \bar{D}$. Due to (5.4) for a small neighborhood U_{x_0} of x_0 we have $\inf_{U_{x_0} \cap \bar{D}} g_j \geq 1/(2K)$ for all j large enough. Let u_0 be the solution of

$$\Delta u_0(x) = N(m, d)(I_{U_{x_0} \cap D}/(2K))^{1-1/m}(x)$$

with zero boundary data, where $N(m, d)$ is the same positive constant as that of (5.3). Then by the comparison principle, we get

$$u_j(y) \leq u_0(y) \quad \text{in } D.$$

However, by the strong maximum principle, u_0 is strictly negative in D and bounded away from 0 on $D \setminus \Delta_{\varepsilon_2}$, which contradicts (5.5).

Now for any $x \in \partial D$, we consider u in the ball B_x . By the previous proof, we have $u \leq -\gamma_0$ on the south half sphere and $u(x) = 0$. Moreover, u is subharmonic in B_x . Denote v to a harmonic function in B_x with boundary data 0 on the north half sphere and $-\gamma_0$ on the south half sphere. By the comparison principle again we have $u(y) \leq v(y)$ in

B_x , and therefore $D_n u(x) \leq D_n v(x) < 0$. Actually, by our construction of v , $D_n v(x)$ only depends on K and d (not x). This completes the proof of the lemma. $\square$

6. INTERIOR SECOND ORDER DERIVATIVES

Here, our goal is to firstly give an interior estimate of the second order derivatives of the solution via the estimates on the boundary of the second order derivatives.

Note that for any function $H(\alpha, x)$ which is twice differentiable in x , if $\inf_{\alpha \in A} H(\alpha, x)$ is also twice differentiable, then for any $\xi \in \mathbb{R}^d$ pointwisely we have

$$\left(\inf_{\alpha \in A} H(\alpha, x) \right)_{(\xi)(\xi)} \leq H_{(\xi)(\xi)}(\alpha_0, x), \quad \Delta \left(\inf_{\alpha \in A} H(\alpha, x) \right) \leq \Delta H(\alpha_0, x),$$

where $\alpha_0 \in A$ such that $H(\alpha_0, x) = \inf_{\alpha \in A} H(\alpha, x)$. After differentiating (4.8) twice in the direction ξ , we get

$$\begin{aligned} & (P_m^{1-1/m} P_{m-1}^{-1})(u_{xx}(x)) \frac{m-1}{d-m+1} [g_{(\xi)(\xi)} g^{-1/m}(x) \\ & - (1/m) g_{(\xi)}^2 g^{-1-1/m}(x)] \leq a^{ij}(u_{xx}(x)) u_{x^i x^j (\xi)(\xi)}(x). \end{aligned}$$

Because of (2.6), (4.1) and (4.6), we obtain

$$P_{m, u_{x^i x^j}}(u_{xx}(x)) u_{x^i x^j (\xi)(\xi)}(x) \geq -N g^{m-2}(x). \quad (6.1)$$

Observe that by Lemma 4.1, (6.1) is an elliptic equation in D . Combining (4.12) with the comparison principle for the elliptic equations in D' , we get an upper estimate

$$u_{(\xi)(\xi)}(x) \leq N + \sup_{\partial D'} u_{(\xi)(\xi)}(x) \leq N + \sup_{\partial D} u_{(\xi)(\xi)}(x). \quad (6.2)$$

The last inequality is because $|u_{(\xi)(\xi)}(x)| \leq d-1$ on $\partial D' \cap D$.

To obtain the lower estimate, it remains to use (4.10) again. As a conclusion, we get

Theorem 6.1. *Let $u \in C^4(D) \cap C^2(\bar{D})$ be the solution of (4.1) in D and satisfies $(u_{xx}(x)) \in C_m$ in D . Then*

$$\sup_D |D^2 u| \leq N(1 + \sup_{\partial D} |D^2 u|), \quad (6.3)$$

where N is a constant depending only on K and d .

Remark 6.2. It turns out that to get (6.3) it suffices to assume Δg to be bounded from below. Indeed, under this condition, instead of (6.1) and (6.2) we have

$$P_{m, u_{x^i x^j}}(u_{xx}(x)) \Delta u_{x^i x^j}(x) \geq -N g^{m-2}(x),$$

$$\Delta u(x) \leq N + \sup_{\partial D'} \Delta u(x) \leq N + \sup_{\partial D} \Delta u(x).$$

The last estimate together with (4.11) yields (6.3).

7. BOUNDARY SECOND ORDER DERIVATIVES

We remark here that for the problem with homogeneous boundary condition the estimate of the second order derivatives on the boundary follows from the arguments in Section 5 of [3]¹ by applying Lemma 5.3 instead of the usual Hopf lemma. Here we give some details for the sake of completeness. While estimating the mixed second derivatives, we use Krylov's approach in [16] and [17].

For any $x \in \partial D$, after a shift of the origin and an orthogonal transformation, we may suppose x is the origin and x^n -axis is the inner normal. By further transforming the coordinate $x' = (x^1, \dots, x^{n-1})$, we can assume in a small neighborhood U_0 of x , ∂D can be represented by $x^n = \bar{\psi}(x')$ and $u_{x_i x_j}(0) = 0$ for $i \neq j$, $i, j = 1, \dots, n-1$. Here $x^n - \bar{\psi}(x')$ is in the class of $C^{3,1}(\bar{U}_0)$ and

$$\|x^n - \bar{\psi}(x')\|_{C^{3,1}(\bar{U}_0)} \leq N(d, K), \quad \nabla \bar{\psi}(0) = 0.$$

Then it suffices to estimate $u_{x^j x^j}(0)$, $u_{x^j x^n}(0)$ and $u_{x^n x^n}(0)$, where $j = 1, \dots, n-1$.

The estimation of the tangential second order derivatives on the boundary is standard (cf. [3], [7] or [17]). We differentiate the equality

$$u(x', \bar{\psi}(x')) = 0$$

twice with respect to x^j , $j = 1, \dots, n-1$, and get

$$u_{x^n}(0) \bar{\psi}_{x^j x^j}(0) + u_{x^j x^j}(0) = 0, \quad (7.1)$$

which along with Theorem 5.2 gives a bound for $u_{x^j x^j}(0)$.

Next, let's estimate the mixed derivatives $u_{x^j x^n}(0)$. We start with introduce a few more objects. Denote $\mathbb{A}$ to be the space of all skew-symmetric matrices and for $p \in \mathbb{A}$ we set

$$a(w, p) = a(e^p w e^{-p}) = e^p a(w) e^{-p}, \quad \sigma = \sqrt{2a}, \quad f(w, p, x) = f(w, x).$$

For $\xi \in \mathbb{R}^d$, we also define $P(w, x)\xi = P(x)\xi$ with value in $\mathbb{A}$ by the formula

$$[P\xi]_{ij} = \psi_{1x^i(\xi)} \psi_{1x^j} - \psi_{1x^i} \psi_{1x^j(\xi)}.$$

Since $e^p C_m e^{-p} = C_m$ and $f(e^p w e^{-p}, x) = f(w, x)$, we have

$$0 = \inf_{w \in C_m} \{a^{ij}(w) u_{x^i x^j}(x) + f(w, x)\}$$

¹We are grateful to the referee for pointing this out.

$$= \inf_{w \in C_m, p \in \mathbb{A}} \{a^{ij}(w, p)u_{x^i x^j}(x) + f(w, p, x)\}.$$

Owing to the proof of Theorem 5.9 of [15], there exist positive numbers δ_1 and δ_2 depending only on K and d such that by taking $B_1 := \delta_1 I_d$ the following assumption is satisfied. This assumption is exactly Assumption 1.2 (d) of [17] with K there equal to 0.

Assumption 7.1. For any $x \in \partial D$, $\xi \perp \psi_{1x}(x)$, $|\xi| = 1$, $w \in C_m$, $p = 0$ we have $(B_1 \xi, \xi) = \delta_1$ and

$$\begin{aligned} L^w \psi_1 + a^{ij} B_{1ij} &\leq -\delta_2 \\ (B_1 \xi, \xi) L^w \psi + \sum_k (\partial(\xi) \psi_{1(\sigma^k)})^2 + 2(B_1 \xi, \sigma^k) \partial(\xi) \psi_{1(\sigma^k)} &\leq -\delta_2, \end{aligned}$$

where

$$\partial(\xi) \psi_{1(\sigma^k)} = \frac{d}{dh} \psi_{1x^i}(x + h\xi) \sigma^{ik}(w, hP(w, x)\xi)|_{h=0}.$$

The estimation of the mixed second order derivatives is a direct application of Theorem 1.10 of [17]. First notice that, as we mentioned before, (4.8) is equivalent to (4.9), which is uniformly elliptic. Next, to estimate the mixed second order derivatives we consider the function $\bar{u} = u/\psi$, which satisfies a higher dimensional elliptic Bellman equation on an auxiliary manifold. Then the problem is reduced to the estimation of tangential first order derivatives of $\bar{u}$ on the manifold. In turn, actually it suffices to have $f(w, \cdot)$ to be in $C^1(\bar{D})$, which is already satisfied in our case due to (4.15). As a conclusion we get

Lemma 7.2. *Under our assumptions, there exist positive constants $\rho = \rho(K, d)$ and $N = N(K, d)$ such that for any $x \in \partial D$ and unit $\tau \perp \psi_x(x)$ we have*

$$|u_{(\tau)(n)}(x)| \leq N(1 + \max_{\partial D(\rho)} (|u| + |u_x|)).$$

This immediately implies the estimate of $u_{x^j x^n}(0)$, $j = 1, \dots, n-1$.

We use the equation (2.4) itself to estimate the normal second order derivative $u_{x^n x^n}(0)$. Equation (2.4) at the origin can be rewritten as

$$u_{x^n x^n}(0) P_{m-1, d-1}(u_{x^1 x^1}(0), \dots, u_{x^{n-1} x^{n-1}}(0)) + G = g^{m-1}(0), \quad (7.2)$$

where G is a sum of products of $u_{x^j x^j}(0)$ and $u_{x^j x^n}(0)$, $j = 1, \dots, n-1$. By the results of Section 7, we have

$$|G| \leq N(K, d). \quad (7.3)$$

Due to (2.1), (7.1) and Lemma 5.3, we get

$$P_{m-1, d-1}(u_{x^1 x^1}(0), \dots, u_{x^{n-1} x^{n-1}}(0))$$

$$\begin{aligned}
 &= (-u_{x^n}(0))^{m-1} P_{m-1,d-1}(\bar{\psi}_{x^1 x^1}(0), \dots, \bar{\psi}_{x^{n-1} x^{n-1}}(0)) \\
 &\geq \delta(D, K, d) P_{m-1,d-1}(\kappa^1, \dots, \kappa^{d-1}) \geq \delta(D, K, d) > 0.
 \end{aligned} \tag{7.4}$$

Combining (7.2), (7.3) and (7.4) together yields

$$|u_{x^n x^n}(0)| \leq N(D, K, d). \tag{7.5}$$

Thus for the non-degenerate case $g > 0$ in D , we get the estimate for $|u_{xx}|$ on ∂D , and subsequently in D by Theorem 6.1. For the general case, we only have to use Lemma 4.6.

Remark 7.3. It is worth noting that when we estimate the second order derivatives on the boundary ∂D , we only use the fact

$$\|f(w, \cdot)\|_{C^1(\bar{D})} \leq N(K, d).$$

So to get the boundary estimates, it suffices to assume $g^{2(m-1)/m} \in C^1(\bar{D})$, which in general is weaker than the condition that g itself is in $C^1(\bar{D})$.

Remark 7.4. Our method can be carried over to a larger class of Hessian equations

$$P_m(u_{xx} + h) = g^{m-1} \tag{7.6}$$

with zero boundary condition, where $h \in C^{1,1}(\bar{D}, \bar{C}_m)$ is a $\bar{C}_m$ -valued function satisfying the following condition:

$$\begin{aligned}
 \text{Tr } h &\leq \frac{md}{d-m+1} \frac{P_m^{1-1/m}(I)}{P_{m-1}(I)} g^{1-1/m}, \\
 \text{Tr } h &\not\equiv \frac{md}{d-m+1} \frac{P_m^{1-1/m}(I)}{P_{m-1}(I)} g^{1-1/m}.
 \end{aligned}$$

Naturally, we look for solutions such that $u_{xx} + h \in \bar{C}_m$ (a.e.).

Indeed, as we mentioned before, for large t on ∂D we have $-\psi_{1,xx} \in C_m$. Due to Corollary 4.2 on ∂D we have $\text{diag}\{\kappa^1, \dots, \kappa^{d-1}\} \in C_{m-1}$. As before, we can rewrite (7.6) as a Bellman equation

$$\inf_{w \in C_m} \{L^w u(x) + f(w, x) + \text{Tr}(a(w)h(x))\} = 0, \tag{7.7}$$

where L^w and $f(w, x)$ are defined in the same way as in (4.8). By the same method we can get the estimates of u and u_x , and reduce the interior estimate of u_{xx} to the estimates of u_{xx} on the boundary.

Under the linear transformation introduced at the beginning of Section 7, $h(0)$ becomes $\bar{h}(0)$. Denote $\bar{h}_n$ to be the $(d-1) \times (d-1)$ submatrix obtained by deleting the n th row and n th column of $\bar{h}(0)$. By Corollary 4.2, $\bar{h}_n$ is in $\bar{C}_{m-1,d-1}$. After estimating the tangential and mixed second order derivatives on the boundary in a similar way,

we can obtain the estimate of $u_{x^n x^n}(0)$ by using the equation (7.6) itself and the inequality

$$\begin{aligned} & P_{m-1,d-1}(\text{diag}\{u_{x^1 x^1}(0), \dots, u_{x^{n-1} x^{n-1}}(0)\} + \bar{h}_n) \\ & \geq P_{m-1,d-1}(\text{diag}\{u_{x^1 x^1}(0), \dots, u_{x^{n-1} x^{n-1}}(0)\}) \\ & = P_{m-1,d-1}(\text{diag}\{-u_{x^n}(0)\bar{\psi}_{x^1 x^1}(0), \dots, -u_{x^n}(0)\bar{\psi}_{x^{n-1} x^{n-1}}(0)\}) \\ & \geq (-u_{x^n}(0))^{m-1} P_{m-1,d-1}(\bar{\psi}_{x^1 x^1}(0), \dots, \bar{\psi}_{x^{n-1} x^{n-1}}(0)) \geq \delta(D, K, d) > 0. \end{aligned}$$

Here in the first step we use the inequality

$$P_{m-1}(A + B) \geq P_{m-1}(A)$$

for any $A, B \in \bar{C}_{m-1}$ (cf. Lemma 4.3 or (4.4)).

8. PROOF OF LEMMA 4.6

Let g_n be a sequence of strictly positive functions in $C^2(\bar{D})$ such that for $n = 1, 2, \dots$, the functions $g_n(x) + 2K|x|^2$ are convex on $\bar{D}$ and

$$\inf_D g_n \geq 1/(2n), \quad \|g_n\|_{C^1(\bar{D})} \leq 2K, \quad \|g_n - g\|_{C^0(\bar{D})} \leq 1/n.$$

By our assumption there exists $N = N(D, d, K)$ such that

$$\|v_n\|_{C^2(\bar{\Omega})} \leq N(D, d, K), \quad m = 1, 2, \dots, \quad (8.1)$$

where $v_n \in C^2(\bar{D}) \cap C^4(D)$ is the solution of (4.1) with g_n in place of g and with zero Dirichlet boundary condition. By Arzelà-Ascoli theorem, after passing to a subsequence if necessary, $\{v_n\}$ converges in $C^1(\bar{D})$ to a function v . Again by (8.1) we get $v \in C^{1,1}(\bar{D})$ and

$$\|v\|_{C^{1,1}(\bar{\Omega})} \leq N(D, d, K).$$

Owing to the uniqueness of the admissible weak solution, we obtain

$$u = v, \quad \|u\|_{C^{1,1}(\bar{\Omega})} \leq N(D, d, K).$$

This completes the proof of (i).

To prove (ii) we use the idea in the proof of Lemma 7.3.4 [14]. Let Ω be a countable dense subset of C_m . Obviously, one has

$$\inf_{w \in C_m} [L^w u + f] = \inf_{w \in \Omega} [L^w u + f].$$

For any $w \in \Omega$, in D it holds that

$$L^w v_n(x) + f_n(w, x) \geq 0, \quad (8.2)$$

where

$$f_n(w, x) = -m(d - m + 1)^{-1} P_m^{1-1/m}(w) P_{m-1}^{-1}(w) g_n^{1-1/m}(x).$$

After multiplying (8.2) by a nonnegative function $\eta \in C_0^\infty(D)$, integrating by parts, passing to the limit over the sequence n and integrating by parts again, we obtain

$$\int_D [a^{ij}(w)\eta(x)u_{x^i x^j}(x) + \eta(x)f(w, x)] dx \geq 0.$$

Because $\eta \in C_0^\infty(D)$ is arbitrary, we further get

$$a^{ij}(w)u_{x^i x^j}(\cdot) + f(w, \cdot) \geq 0 \quad \text{a.e. in } D.$$

Since Ω is a countable set, we reach

$$\sup_{w \in \Omega} [Lu(x) + f(w, x)] \geq 0 \quad \text{a.e. in } D. \quad (8.3)$$

Next we prove the opposite inequality. Here we use again the method by which Lemma 7.3.4 of [14] is proved. Let $\varepsilon < 1$ be a positive number. Recall that v_n satisfies the Bellman equation

$$\inf_{w \in C_m} \{L^w v_n(x) + f_n(w, x)\} = 0.$$

Given any n such that $1/n < \varepsilon$, we have

$$\begin{aligned} & \inf_{w \in C_m} \{L^w v_n(x) + \varepsilon \Delta v_n(x) + f(w, x)\} \\ & \leq \sup_{w \in C_m} \{\varepsilon \Delta v_n(x) + f(w, x) - f_n(w, x)\} \leq N(D, d, K)\varepsilon, \\ & \inf_{w \in C_m} \{L^w v_n(x) + \varepsilon \Delta v_n(x) + f(w, x) - N(D, d, K)\varepsilon\} \leq 0. \end{aligned}$$

Note that the elliptic operator $\inf_{w \in C_m} [L^w + \varepsilon \Delta]$ is uniformly non-degenerate. Owing to Theorem 3.6.3 [14], after passing to the limit over the sequence n we obtain

$$\inf_{w \in C_m} \{L^w u(x) + \varepsilon \Delta u(x) + f(w, x) - N(D, d, K)\varepsilon\} \leq 0 \quad \text{a.e. in } D.$$

Letting $\varepsilon \downarrow 0$ yields an inequality opposite to (8.3). The lemma is proved.

REFERENCES

- [1] L.A. Caffarelli, L. Nirenberg, J. Spruck, The Dirichlet problem for nonlinear second-order elliptic equations, I: Monge-Ampère equation, *Comm. Pure Appl. Math.*, **37** (1984), 369–402.
- [2] L.A. Caffarelli, J.J. Kohn, L. Nirenberg, J. Spruck, The Dirichlet problem for nonlinear second-order elliptic equations, II: Complex Monge-Ampère equation and uniformly elliptic equations, *Comm. Pure Appl. Math.*, **38** (1985), 209–252.
- [3] L.A. Caffarelli, L. Nirenberg, J. Spruck, The Dirichlet problem for nonlinear second-order elliptic equations, III. Functions of the eigenvalues of the Hessian, *Acta Math.*, **155** (1985), 261–301.
- [4] L.A. Caffarelli, L. Nirenberg, J. Spruck, The Dirichlet problem for degenerate Monge-Ampère equations. *Rev. Mat. Iberoamericana*, **2** (1986), 19–27.

- [5] L. Garding, An inequality for hyperbolic polynomials, *J. Math. Mech.*, **8** (1959), 957–965.
- [6] G. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, 2nd edition, Springer-Verlag, Berlin-New York, 1983.
- [7] P. Guan, N.S. Trudinger, X. Wang, On the Dirichlet problem for degenerate Monge-Ampère equations, *Acta Math.*, **182** (1999), no.1, 87–104.
- [8] L. Hörmander, *The analysis of linear partial differential operators*, Vols. 1,2, Springer, New York, 1990.
- [9] J. Hong, The estimates near the boundary for solutions of Monge-Ampère equations. *J. Partial Differential Equations*, **7** (1994), 97–110.
- [10] N.M. Ivochkina, A priori estimate of $\|u\|_{C^2(\bar{\Omega})}$ of convex solutions of the Dirichlet problem for the Monge-Ampère equation, *Zap. Nauchn. Sem. Leningrad. otдел. Mat. Inst. Steklov.* **96** (1980), 69–79 (Russian); English transl. *J. Soviet Math.*, **21** (1983), 689–697.
- [11] N. Ivochkina, N.S. Trudinger, X. Wang, The Dirichlet problem for degenerate Hessian equations, *Comm. Partial Differential Equations*, **29** (2004), No. 1-2, 219–235.
- [12] N.V. Krylov, *Controlled diffusion processes*, Springer-Verlag, New York, 1980.
- [13] N.V. Krylov, Boundedly inhomogeneous elliptic and parabolic equations in a domain, *Izv. Akad. Nauk SSSR*, **47** (1983), 75–108 (Russian); *English transl. Math. USSR-Izv.*, **22** (1984), 67–98.
- [14] N.V. Krylov, *Nonlinear elliptic and parabolic equations of second order*, Nauka, Moscow, 1985; English transl., Reidel, Dordrecht, 1987.
- [15] N.V. Krylov, On the general notion of fully nonlinear second order elliptic equation, *Trans. Amer. Math. Soc.*, **347** No. 3, (1995), 857–895.
- [16] N.V. Krylov, Weak interior second order derivative estimates for degenerate nonlinear elliptic equations, *Differential and Integral Equations*, **7** (1994), 133–156.
- [17] N.V. Krylov, A theorem on degenerate elliptic Bellman equations in bounded domains, *Differential and Integral Equations*, **8** (1995), 961–980.
- [18] N.V. Krylov, Smoothness of the payoff function for a controllable process in a domain, *Izv. Akad. Nauk SSSR, ser. math.*, **53** (1989), 66–96, in Russian; English translation: *Math. USSR-Izv.*, **34**, (1990), 65–95.
- [19] Pogorelov, A., *The Minkowski multidimensional problems*, J. Wiley, New York, 1978.
- [20] M.V. Safonov, The classical solution of Bellman’s elliptic equation, *Dokl. Akad. Nauk SSSR*, **278**, No. 4 (1984), 810–813; English transl., *Soviet Math. Dokl.*, **30** (1984), No. 42, 482–485.
- [21] N.S. Trudinger, On the Dirichlet problems for Hessian equations, *Acta Math.*, **175** (1995), 151–164.
- [22] N.S. Trudinger, Weak solutions of Hessian equations, *Comm. Partial Differential Equations*, **22** (1997), no. 7-8, 1251–1261.
- [23] X. Wang, Some counterexamples to the regularity of Monge-Ampère equations, *Proc. Amer. Math. Soc.*, **123** (1995), 841–845.

(H. Dong) 127 VINCENT HALL, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455, USA

E-mail address: `hjdong@math.umn.edu`